



THE UNIVERSITY OF TEXAS AT AUSTIN

ANIMAL RESOURCES CENTER

2701 Speedway · Austin, TX 78712-0136

(512) 471-7534 · FAX (512) 471-4336 · www.utexas.edu/research/arc

Hamster-Specific Anesthesia

I. Purpose:

This document has been designed by the ARC veterinary staff as guidance for hamster anesthesia. This is not intended to be an inclusive list of all possible drug combinations that can be used in hamsters. Instead, these guidelines are general recommendations. Consequently, they do not include reference to specific research-associated concerns. If you have questions about the use of anesthetics for your particular situation, please work with your area veterinarian to develop the most effective anesthetic plan.

Anesthesia is the loss of feeling in all or part of the body, with or without loss of consciousness. Animals may be anesthetized for surgery, for non-surgical procedures that may be painful, or for non-painful procedures that require immobilization. Steps must be taken before, during, and after anesthesia to ensure the safety of the animal and efficacy of anesthesia. These are listed in the General Guidelines section below.

Anesthetic drugs can be administered parenterally or by inhalation. Commonly used anesthetic agents are described below. The choice of anesthetic agent will depend on the procedure to be performed, research aims, and other factors. Consult your area veterinarian with questions about drug selection.

II. General Guidelines and Considerations for Hamster Anesthesia:

- **Acclimation:** A one-week post-shipping acclimation period is recommended for all animals to prevent stress-induced responses. At a minimum, animals require 72 hours to regain basic physiological baselines.
- **Fasting:** Pre-anesthetic fasting is usually not necessary for hamsters due to their inability to vomit. However, if fasting is employed (e.g., to stabilize body weights before anesthesia) limit to no more than 2-3 hours due to the high metabolic rate of small rodents. **Never restrict water.**

- Eye protection: Hamster eyes remain open under anesthesia. This can lead to corneal drying and trauma. Apply ophthalmic ointment (e.g., Paralube® or Lacrilube®) to eyes if:
 - The anesthetic event lasts longer than 5 minutes.
 - Anesthesia is being delivered by facemask.

- Heat support: All species are at risk for hypothermia while under anesthesia. Hamsters are particularly susceptible due to their high body surface area to body mass ratio. Hypothermia induces a significant physiological stress on animals that can prolong recovery and potentially be fatal.
 - Provide supplemental heat during all anesthetic events.
 - Supplemental heat sources include circulating water blankets, electric heating pads, and commercial products that can be warmed in a microwave or water bath (such as isothermic pads or gel warmers).
 - Never place the animal directly on the heat source.
 - Contact your area veterinarian with questions about using or purchasing supplemental heat sources.

- Fluid support: Consider providing warm SQ or IP fluids, particularly for prolonged anesthetic events or animals that are ill, aged, or debilitated. Consult your veterinarian for additional information.

- Monitoring: Regardless of the anesthetic administered, the hamster must be monitored to avoid excessive depression of cardiac and respiratory functions, or insufficient anesthesia.

- Parameters that can be monitored in an anesthetized hamster without specialized equipment include:
 - Anesthetic depth- toe pinch
 - Respiratory rate and pattern (normal undisturbed rate = ~35-135/min, a slow rate drop of 50% is acceptable during anesthesia)
 - Mucous membrane color (should be pink nt blue or grey).
 - Oxygen saturation (>95%) and heart rate (350-500 beats/min) can be monitored using a pulse oximeter.

- Recovery: Continue to monitor animals until they are fully recovered.
 - Recover animals on paper towels (without bedding) in a clean cage. This minimizes the risk of tracheal obstruction or pneumonia.
 - Recover anesthetized animals alone in a cage.
 - Continue to provide supplemental heat during recovery.
 - When the animal is ambulatory, return it to the home cage with immediate access to food and water.
 - Depending on the surgery performed, hamsters may need to be housed individually in the post-operative period to avoid suture gnawing or incision-site trauma caused by cagemates. Any plans for individual housing should be included in the IACUC protocol so that they can be reviewed and approved.

III. Anesthetic Drugs and Procedures:

1. **Inhaled Anesthetics:** Isoflurane is the preferred inhaled anesthetic. It has rapid and reliable onset and recovery.

A. Inhalation anesthesia may be delivered by the drop jar method, or by induction chamber, facemask, or endotracheal tube using a precision vaporizer.

i. Drop jar:

- a. Do not allow the hamster to come into direct contact with the liquid inhalant anesthetic (which can be achieved by placing an impermeable mesh grid over the cotton/gauze).
- b. Because the anesthetic concentration cannot be controlled in the chamber and can rapidly reach toxic concentrations, this method is reserved for very short procedures such as tail biopsies, subcutaneous tumor implantation, or induction prior to facemask anesthesia.

ii. Induction chambers, facemasks and endotracheal tubes:

- a. These require gas anesthesia machines with an oxygen source and a precision vaporizer. Some research units may use room air rather than pure oxygen as a gas source which may also be acceptable.
- b. The vaporizer must be compatible with the specific inhaled anesthetic agent.
- c. Due to the small respiratory capacity in hamsters, use a non-rebreathing system.

iii. When using inhalant anesthesia, use a fume hood or an anesthetic system equipped with a gas scavenging system to minimize occupational exposure to exiting gases.

Drug	Dose	Oxygen flow rate	Comments
Isoflurane Recommended	4-5% for induction 1-2% for maintenance	300-600 ml/min	Approximately 0.6 mls of liquid isoflurane per liter of chamber volume

B. **Injectable Anesthetics:** Combined ketamine/xylazine is the preferred injectable anesthetic in hamsters. Other commonly used injectable agents are listed below.

i. Hamsters vary significantly in their sensitivity to various anesthetics. Age, body composition, strain, health status, genetic manipulation, and sex are just a few of the factors that can contribute to anesthetic sensitivity. The following doses are general guidelines that may vary significantly based on the aforementioned factors. When adding anesthetics to your protocol, it is advisable to provide a range to allow titration for the specific needs of each animal.

- ii. Intramuscular (IM) injections are not recommended in hamsters as complications such as tissue irritation, lameness and self-mutilation can result.

Injectable Anesthetics:

Drug	Dose	Route	Duration of Anesthesia	Comments
Ketamine + Xylazine *	50-200 mg/kg Ket + 5-10 mg/kg Xyl.	IP	30-60 minutes	Thermal support is crucial. To prolong anesthesia, supplement with 1/3 dose of ketamine only. Xylazine can be reversed with 1 - 2 mg/kg yohimbine IP or a volume of atipamezole equal to the volume of Xylazine (20 mg/ml) given (IP)**.
Ketamine + Xylazine + Acepromazine	50-100 mg/kg Ket + 10 mg/kg Xyl + 1 mg/kg Ace	IP	45-120 mins	Acepromazine prolongs the duration of anesthesia. Thermal support is crucial. If more anesthetic time is needed, supplement with 1/3 dose Ketamine alone. Xylazine can be reversed with 1 - 2 mg/kg yohimbine IP or a volume of atipamezole equal to the volume of Xylazine (20 mg/ml) given (IP)**.
Ketamine +Dexmedetomidine	50-75 mg/kg Ket + 0.25 mg/kg Dexmed	IP	30-60 mins	Thermal support is crucial. If additional anesthetic is needed, supplement with 1/3 dose of ketamine only. Do not re-dose Dexmedetomidine. Dexmedetomidine can be reversed a volume of atipamezole equal to the volume of Dexmedetomidine given (IP)**

* A good starting point is 100 mg/kg Ketamine and 7 mg/kg Xylazine

**Atipamazole (5 mg/ml) is 10X the concentration of Dexmedetomidine (0.5 mg/ml). You need 4-6X as many mg of Atipamaozle (5 mg/ml) to reverse Dexmedetomidine. However, it also has a very high safety margin, so giving an equal volume of Atipamazole to Dexmedetomidine gives you 10 X as many mg of Atipamazole, it is safe, and it allows for reversal without calculating a dose if reversal is urgent.. Equal volumes of Atipamaozle (5 mg/ml) to reverse small animal Xylazine (20 mg/ml) also works because even though Xylazine has a stronger concentration (20 mg/ml), Atipamazole (5 mg/ml) is

cleaner and binds more tightly to the alpha 2 receptor, so equal volumes allows for reversal of xylazine (20 mg/ml) as well.

References:

1. Carpenter JW: Exotic Animal Formulary 3rd edition. Elsevier Saunders 2005.
2. Joanne Sohn and Marcelo A. Couto. Chapter 8 in The Laboratory Rabbit, Guinea Pig, Hamster, and Other Rodents. 2012, Pages 195-215.
3. Laboratory Animal Medicine, 2nd edition. Fox, JG et al. editors. Academic Press, 2002.
4. Wixson, SK and Smiler KL: Anesthesia and analgesia in rodents. In: Anesthesia and analgesia in laboratory animals. Kohn, DF et al. editors. Academic Press, 1997.
5. ULAM Hamster Anesthesia and Analgesia Guidelines.
<https://files.umms.med.umich.edu/ULAM/SOPs/Veterinary/Anesthesia%20Guidelines/ULAM%20SOP%20Hamster%20Anesthesia.pdf>.